

ARITHMETIC GROUP SYMMETRY AND FINITENESS PROPERTIES OF TORELLI GROUPS

ALEXANDRU DIMCA¹ AND STEFAN PAPADIMA²

ABSTRACT. We examine groups whose resonance varieties, characteristic varieties and Sigma-invariants have a natural arithmetic group symmetry, and we explore implications on various finiteness properties of subgroups. We compute resonance varieties, characteristic varieties and Alexander polynomials of Torelli groups, and we show that all subgroups containing the Johnson kernel have finite first Betti number, when the genus is at least 4. We also prove that, in this range, the I -adic completion of the Alexander invariant is finite-dimensional, and the Kahler property for the Torelli group implies the finite generation of the Johnson kernel.

CONTENTS

1. Introduction and statement of results	1
2. Associated graded Lie algebra	5
3. Resonance varieties of Torelli groups	7
4. Resonance and finiteness properties	10
5. Characteristic varieties of Torelli groups	14
6. Sigma-invariants and the Kahler property	18
References	22

1. INTRODUCTION AND STATEMENT OF RESULTS

The natural $\text{Aut}(T)$ -symmetry factors through $\text{Out}(T)$, for a good number of invariants of a group T . The simplest examples are the abelianization, T_{ab} , and the quotient of T_{ab} by its torsion, T_{abf} . More well-known examples are provided by the

2000 *Mathematics Subject Classification*. Primary 20J05, 57N05; Secondary 11F23, 20G05.

Key words and phrases. resonance variety, characteristic variety, Alexander invariant, Alexander polynomial, Sigma-invariant, arithmetic group, semisimple Lie algebra, finiteness properties, Torelli group, Johnson kernel, Kahler group.

¹ Partially supported by the French-Romanian Programme LEA Math-Mode and ANR-08-BLAN-0317-02 (SEDIGA).

² Partially supported by the French-Romanian Programme LEA Math-Mode and CNCSIS-UEFISCSU project PNII-IDEI 1189/2008.

cohomology ring $H^\bullet T := H^\bullet(T, \mathbb{C})$ and the graded Lie algebra associated to the lower central series, $\mathcal{G}_\bullet(T) := \text{gr}_\bullet T \otimes \mathbb{C}$.

Our starting point in this work is to consider a group epimorphism, $p : \Gamma \twoheadrightarrow D$, with finitely generated kernel T , and to examine three other known types of invariants with natural outer symmetry, through the prism of the D -symmetry induced by the canonical homomorphism, $D \rightarrow \text{Out}(T)$. Firstly, we look at the resonance varieties $\mathcal{R}_k^i(T)$ (i.e., the jump loci for a certain kind of homology, associated to the ring $H^\bullet T$), sitting inside $H^1 T$; they are reviewed in Section 3, and their outer symmetry is discussed in Remark 3.1. Secondly, we inspect the characteristic varieties $\mathcal{V}_k^i(T)$ (i.e., the jump loci for homology with rank one complex local systems), lying inside the character torus $\mathbb{T}(T) := \text{Hom}(T_{\text{ab}}, \mathbb{C}^*)$, and their intersection with $\mathbb{T}^0(T) := \text{Hom}(T_{\text{abf}}, \mathbb{C}^*)$; their definition is recalled in Section 5, and their outer symmetry is explained in Lemma 5.1. Finally, we recollect in Section 6 a couple of relevant facts about the Bieri-Neumann-Strebel-Renz (BNSR) invariants, $\Sigma^q(T, \mathbb{Z}) \subseteq H^1(T, \mathbb{R}) \setminus \{0\}$, and we point out their outer symmetry in Lemma 6.1.

Our choice for the types of invariants was dictated by the fact that each of them controls a certain kind of finiteness properties. We begin with resonance, for which this relationship seems to be new. We devote Section 4 to the analysis of the connexions between triviality of resonance and finiteness of various (completed) Alexander-type invariants, for finitely generated groups.

Our motivation comes from the Torelli groups, T_g , consisting of the isotopy classes of homeomorphisms of a closed, oriented, genus g surface, inducing the identity on the first $\mathbb{Z}$ -homology group of the surface. Their finite generation was proved by D. Johnson [21], for $g \geq 3$. According to Hain [16], T_g is a 1-formal group in the sense of D. Sullivan [36], for $g \geq 6$, but T_3 is not 1-formal.

Note also that the full mapping class group Γ_g is an extension of the integral symplectic group $Sp_g(\mathbb{Z})$ by T_g , for $g \geq 1$:

$$(1.1) \quad 1 \rightarrow T_g \rightarrow \Gamma_g \rightarrow Sp_g(\mathbb{Z}) \rightarrow 1.$$

In particular, the resonance and characteristic varieties of T_g , as well as its BNSR-invariants, acquire a natural $Sp_g(\mathbb{Z})$ -symmetry, for $g \geq 3$.

Let $K \subseteq T$ be a subgroup containing the derived group T' . Then $H_1 K := H_1(K, \mathbb{C})$ becomes in a natural way a module over the group ring $\mathbb{C}T_{\text{ab}}$, with module structure induced by T -conjugation. Its I -adic completion is denoted $\widehat{H_1 K}$, where $I \subseteq \mathbb{C}T_{\text{ab}}$ is the augmentation ideal. When $K = T'$, $H_1 K$ is the classical Alexander invariant from link theory (over $\mathbb{C}$). The technique of I -adic completion was promoted in low-dimensional topology by Massey [28]. Here is our first main result.

Theorem A. *Let T be a finitely generated group.*

- (1) *Assume T is 1-formal. Then $\mathcal{R}_1^1(T) \subseteq \{0\}$ if and only if $\dim_{\mathbb{C}} \widehat{H_1 T'} < \infty$.*

- (2) Let $K \subseteq T$ be a subgroup containing T' . Then $\dim_{\mathbb{C}} \widehat{H_1 K} < \infty$, if $\mathcal{R}_1^1(T) \subseteq \{0\}$.

Aiming at finer finiteness properties, we go on by examining characteristic varieties, in Section 5. Here, we start from a basic result of Dwyer and Fried [14], as refined in [31]. It says that the finiteness of Betti numbers, up to degree q , of normal subgroups with free abelian quotient is detected precisely by the finiteness of the intersection between characteristic varieties of type $\mathcal{V}_1^i$, for $i \leq q$, and the corresponding connected subtorus of $\mathbb{T}^0$.

Given the D -symmetry of characteristic varieties, we are thus led to consider the following context, inspired by Torelli groups. Let L be a D -module which is finitely generated and free as an abelian group. Assume that D is an arithmetic subgroup of a simple $\mathbb{C}$ -linear algebraic group S defined over $\mathbb{Q}$, with $\mathbb{Q} - \text{rank}(S) \geq 1$. Suppose also that the D -action on L extends to an irreducible, rational S -representation in $L \otimes \mathbb{C}$. (Note that the above assumptions are satisfied for $g \geq 3$ by $D = Sp_g(\mathbb{Z}) \subseteq Sp_g(\mathbb{C}) = S$ and $L = (T_g)_{\text{abf}}$, due to Johnson's pioneering results on the symplectic symmetry of Torelli groups from [20, 23].) The D -representation in L gives rise to a natural D -action on the connected affine torus $\mathbb{T}(L) := \text{Hom}(L, \mathbb{C}^*)$.

Theorem B. *If the D -module L satisfies the above assumptions, then $\mathbb{T}(L)$ is geometrically D -irreducible, that is, the only D -invariant, Zariski closed subsets of $\mathbb{T}(L)$ are either equal to $\mathbb{T}(L)$, or finite.*

We deduce Theorem B from a deep result in diophantine geometry, due to M. Laurent [25], in Section 5. Note that the conclusion of our theorem above is in marked contrast with the behavior of the induced D -representation in the affine space $L \otimes \mathbb{C}$, for which S -invariant, infinite and proper Zariski closed subsets may well exist. Theorem B enables us to obtain in Section 5 the following consequences of the triviality of resonance.

Theorem C. *Let $p : \Gamma \twoheadrightarrow D$ be a group epimorphism with finitely generated kernel T , having the property that $\mathcal{R}_1^1(T) \subseteq \{0\}$. Assume $D \subseteq S$ is arithmetic, where the $\mathbb{C}$ -linear algebraic group S is defined over $\mathbb{Q}$, simple, with $\mathbb{Q} - \text{rank}(S) \geq 1$. Suppose moreover that the canonical D -representation in T_{abf} extends to an irreducible, rational S -representation in $T_{\text{abf}} \otimes \mathbb{C}$. The following hold.*

- (1) *The intersection $\mathcal{V}_k^1(T) \cap \mathbb{T}^0(T)$ is finite, for all $k \geq 1$.*
- (2) *If moreover $b_1(T) > 1$, the Alexander polynomial Δ^T is a non-zero constant, modulo the units of the group ring $\mathbb{Z}T_{\text{abf}}$.*
- (3) *For any subgroup $N \subseteq T$, containing the kernel of the canonical map, $T \twoheadrightarrow T_{\text{abf}}$, the first Betti number $b_1(N)$ is finite.*

Note that the computation of characteristic varieties and Alexander polynomials can be a very difficult task, in general. What makes life easier in Theorem C, Parts

(1) and (2), is the arithmetic symmetry. As far as Torelli groups are concerned, we obtain the following results. Denote by K_g the kernel of the canonical map, $T_g \twoheadrightarrow (T_g)_{\text{abf}}$, identified by Johnson in geometric terms in [22].

Theorem D. *Let T_g be the Torelli group in genus $g \geq 4$. The following hold.*

- (1) *The resonance (in degree 1 and depth 1) is trivial: $\mathcal{R}_1^1(T_g) = \{0\}$.*
- (2) *The (reduced) characteristic varieties $\mathcal{V}_k^1(T_g) \cap \mathbb{T}^0(T_g)$ are finite, for all $k \geq 1$.*
- (3) *The Alexander polynomial Δ^{T_g} is a non-zero constant, modulo units.*
- (4) *For any subgroup $N \subseteq T_g$ containing the Johnson kernel K_g , the vector space $H_1(N, \mathbb{C})$ is finite-dimensional.*
- (5) *For any subgroup $K \subseteq T_g$ containing T'_g , $\dim_{\mathbb{C}} \widehat{H_1 K} < \infty$.*

We compute resonance varieties of Torelli groups in Section 3. Note that $\mathcal{R}_1^1(T_3) = H^1 T_3$ is non-trivial; this is an easy consequence of a basic result of Hain [16], who computed the graded Lie algebra $\mathcal{G}_{\bullet}(T_g)$, truncated up to degree $\bullet = 2$, for $g \geq 3$. His result, formulated in the category of $Sp_g(\mathbb{C})$ -representations, is recorded in Section 2. We obtain the triviality of $\mathcal{R}_1^1(T_g)$, for $g \geq 4$, by exploiting the $Sp_g(\mathbb{C})$ -symmetry. Parts (2)-(3) of Theorem D follow from Theorem C.

Parts (4)-(5) of Theorem D appear to be rather surprising, since the general belief in the literature seems to be that Torelli groups and other related groups should exhibit *infiniteness* properties; see for instance [15, §5.1] and the survey [17]. Theorem D(4) is a consequence of Theorem C(3), and Theorem D(5) follows from Theorem A(2). Note that the finite-dimensionality of the (uncompleted) Alexander invariant $H_1(T'_g, \mathbb{C})$ cannot be deduced from Theorem D(2), since $(T_g)_{\text{ab}}$ contains non-trivial 2-torsion, according to Johnson [23]. To the best of our knowledge, the finiteness of $b_1(T'_g)$ is an open question.

Theorem D(4) solves two of the most popular finiteness problems related to Torelli groups, as recorded in Farb's list from [15]: for $N = K_g$, we answer Question 5.2 (over $\mathbb{C}$); the general case ($K_g \subseteq N \subseteq T_g$) settles Problem 5.3 (at the level of first Betti numbers).

According to [6, erratum], the finite generation of K_g is an open question, for $g \geq 3$. Another important problem concerns the Kahler property: is T_g a Kahler group (that is, the fundamental group of a compact Kahler manifold)? The answer is negative, in low genus: T_2 is not finitely generated, as proved by McCullough and Miller [29], and T_3 violates the 1-formality property of Kahler groups, established by Deligne, Griffiths, Morgan and Sullivan in [10]. Theorem 4.9 from [15] states that T_g is not a Kahler group for $g \geq 2$. Since the proof assumed infinite generation of K_g , the Kahler group problem is open as well.

We investigate in Section 6 the BNSR invariants, which control the finiteness properties of normal subgroups with abelian quotient [4, 5]. We use their arithmetic symmetry, together with Delzant's description from [11] of the first BNSR invariant

of Kahler groups, and our Theorem D(1), to prove the next result. It connects the Kahler problem for Torelli groups with the finite generation question for Johnson kernels. See Remark 6.6 for more details on our strategy of proof.

Theorem E. *If T_g is a Kahler group, the Johnson kernel K_g must be finitely generated, for $g \geq 4$.*

2. ASSOCIATED GRADED LIE ALGEBRA

The symplectic symmetry is well-known to be an important tool for the study of Torelli groups. In this section, we review a basic result from Hain's work [16], related to the symplectic symmetry at the level of the associated graded Lie algebra.

Let Σ_g be a closed, oriented, genus $g \geq 1$ surface. Denote by Γ_g the associated *mapping class group*, i.e., the group of isotopy classes of orientation-preserving homeomorphisms of Σ_g . Let $Sp_g(\mathbb{Z})$ be the group of symplectic automorphisms of $H := H_1(\Sigma_g, \mathbb{Z})$. The arithmetic group $Sp_g(\mathbb{Z})$ is a Zariski dense subgroup of the semisimple algebraic group $Sp_g(\mathbb{C})$; see e.g. [33, Corollary 5.16]. The *Torelli group* T_g is the kernel of the natural homomorphism p , which associates to an element of Γ_g the induced action on H . The defining exact sequence of the group T_g is

$$(2.1) \quad 1 \rightarrow T_g \rightarrow \Gamma_g \xrightarrow{p} Sp_g(\mathbb{Z}) \rightarrow 1.$$

R. Hain's starting point is D. Johnson's pioneering work from [20, 21, 22, 23], which we review first. In the sequel we will assume that $g \geq 3$. In this range, Johnson proved [21] that the group T_g is finitely generated. For a group G , we denote by G_{abf} the quotient of its abelianization by the torsion subgroup. Among other things, Johnson [20, 23] gave a very convenient description of $(T_g)_{\text{abf}}$, in the following way.

Fix a symplectic basis of H , $\{a_1, \dots, a_g, b_1, \dots, b_g\}$, and denote by $\omega = \sum_{i=1}^g a_i \wedge b_i \in \bigwedge^2 H$ the symplectic form. Note that the $Sp_g(\mathbb{Z})$ -action on H canonically extends to a $Sp_g(\mathbb{Z})$ -action on the exterior algebra $\bigwedge^* H$, by graded algebra automorphisms. Consider the $Sp_g(\mathbb{Z})$ -equivariant embedding, $H \hookrightarrow \bigwedge^3 H$, given by $h \in H \mapsto h \wedge \omega \in \bigwedge^3 H$, and denote by L the $Sp_g(\mathbb{Z})$ -module $\bigwedge^3 H / H$. Johnson's homomorphism, constructed in [20], will be denoted by $J: T_g \rightarrow L$.

Theorem 2.1 (Johnson). *The group homomorphism J is Γ_g -equivariant, with respect to the (left) conjugation action on T_g induced by (2.1), and the restriction of the $Sp_g(\mathbb{Z})$ -action on L via p . It induces a $Sp_g(\mathbb{Z})$ -equivariant isomorphism, $J: (T_g)_{\text{abf}} \xrightarrow{\cong} L$.*

Setting $H(\mathbb{C}) = H \otimes \mathbb{C}$ and $L(\mathbb{C}) = L \otimes \mathbb{C}$, it follows that the canonical representation of $Sp_g(\mathbb{Z})$, coming from (2.1), in $(T_g)_{\text{ab}} \otimes \mathbb{C} \simeq L(\mathbb{C})$, extends to a rational representation of $Sp_g(\mathbb{C})$. This symplectic symmetry propagates to higher degrees, in the following sense.

Recall that the *associated graded Lie algebra* (with respect to the lower central series) of a group G , $\mathrm{gr}_\bullet G$, is generated as a Lie algebra by $\mathrm{gr}_1 G = G_{\mathrm{ab}}$.

Lemma 2.2. *Given a group extension,*

$$(2.2) \quad 1 \rightarrow T \rightarrow \Gamma \rightarrow D \rightarrow 1,$$

assume that T is finitely generated, D is a Zariski dense subgroup of a complex linear algebraic group S , and the D -action on T_{ab} extends to a rational representation of S in $T_{\mathrm{ab}} \otimes \mathbb{C}$. Then the D -action on $\mathrm{gr}_\bullet T$ extends to an action of S on $\mathrm{gr}_\bullet T \otimes \mathbb{C}$, in the category of graded rational representations; moreover, every $s \in S$ acts on $\mathrm{gr}_\bullet T \otimes \mathbb{C}$ by a graded Lie algebra automorphism.

Proof. Presumably this result is well-known to the experts. Being unable to find a reference, we decided to include a proof. Set $\mathcal{T}_\bullet := \mathrm{gr}_\bullet T \otimes \mathbb{C}$, noting that D acts on $\mathcal{T}_\bullet$ by graded Lie algebra automorphisms. For each $q \geq 1$, denote by $K_q \subseteq \mathcal{T}_1^{\otimes q}$ the kernel of the linear surjection sending $t_1 \otimes \cdots \otimes t_q$ to $\mathrm{ad}_{t_1} \circ \cdots \circ \mathrm{ad}_{t_{q-1}}(t_q) \in \mathcal{T}_q$. Since $D \subseteq S$ is Zariski dense, the linear subspace K_q is S -invariant. It remains to show that the rational representations of S in $\mathcal{T}_\bullet$ constructed in this way, which extend the D -action coming from (2.2), have the property that $s[a, b] = [sa, sb]$, for $s \in S$, $a \in \mathcal{T}_q$ and $b \in \mathcal{T}_r$. This in turn is easily proved by induction on q . Induction starts with $q = 1$, by noting that the iterated Lie bracket, $\mathcal{T}_1^{\otimes r+1} \rightarrow \mathcal{T}_{r+1}$, is S -equivariant by construction. For the inductive step, write $a = [t, a']$, with $t \in \mathcal{T}_1$ and $a' \in \mathcal{T}_{q-1}$, and use the Jacobi identity to conclude. $\square$

By Lemma 2.2 and Theorem 2.1, we have a short exact sequence of rational $Sp_g(\mathbb{C})$ -representations,

$$(2.3) \quad 0 \rightarrow \mathcal{K} \rightarrow \bigwedge^2 \mathrm{gr}_1 T_g \otimes \mathbb{C} \xrightarrow{\beta} \mathrm{gr}_2 T_g \otimes \mathbb{C} \rightarrow 0,$$

where β denotes the Lie bracket. Hain [16] computed the associated exact sequence of $\mathfrak{sp}_g$ -modules, where $\mathfrak{sp}_g$ is the Lie algebra of $Sp_g(\mathbb{C})$. To describe his result, we follow the conventions from [16, Section 6]. Our references for algebraic groups (respectively Lie algebras) are [19] (respectively [18]).

The Lie algebra of the maximal diagonal torus in $Sp_g(\mathbb{C})$ is denoted $\mathfrak{h}$, and has coordinates $t = (t_1, \dots, t_g)$. Let $\Phi \subseteq \mathfrak{h}^*$ be the corresponding root system, with the standard choice of positive roots, $\Phi^+ = \{t_i - t_j, t_i + t_j \mid 1 \leq i < j \leq g\} \cup \{2t_i \mid 1 \leq i \leq g\}$. Let $\mathfrak{sp}_g := \mathfrak{s} = \mathfrak{n}^- \oplus \mathfrak{h} \oplus \mathfrak{n}^+$ be the canonical decomposition of the Lie algebra. We denote by B the associated Borel subgroup of $Sp_g(\mathbb{C}) := S$, with unipotent radical U ; the Lie algebra of B is $\mathfrak{h} \oplus \mathfrak{n}^+$, and the Lie algebra of U is $\mathfrak{n}^+$. We work with the finite-dimensional $\mathfrak{s}$ -modules associated to rational representations of S . The irreducible ones are of the form $V(\lambda)$, where the dominant weight λ is a positive integral linear combination of the fundamental weights, $\{\lambda_j(t) = t_1 + \cdots + t_j \mid 1 \leq j \leq g\}$.

It follows from Theorem 2.1 that $\mathrm{gr}_1 T_g \otimes \mathbb{C} = V(\lambda_3)$, as $\mathfrak{sp}_g$ -modules. According to [16, Lemma 10.2], all irreducible submodules of $\bigwedge^2 V(\lambda_3)$ occur with multiplicity one, and $\bigwedge^2 V(\lambda_3)$ contains $V(2\lambda_2) \oplus V(0)$ as a submodule.

Theorem 2.3 (Hain). *The $\mathfrak{sp}_g$ -map β from (2.3) is the canonical $\mathfrak{sp}_g$ -equivariant projection of $\bigwedge^2 V(\lambda_3)$ onto the submodule $V(2\lambda_2) \oplus V(0)$.*

3. RESONANCE VARIETIES OF TORELLI GROUPS

In this section, we use representation theory to compute the resonance varieties (in degree 1 and for depth 1) of the Torelli groups T_g , for $g \geq 3$, over $\mathbb{C}$.

We begin by reviewing the *resonance varieties*, $\mathcal{R}_d^i(A^\bullet)$, associated to a connected, graded-commutative $\mathbb{C}$ -algebra $A^\bullet$, for (degree) $i \geq 0$ and (depth) $d \geq 1$. Given $a \in A^1$, denote by μ_a left-multiplication by a in $A^\bullet$, noting that $\mu_a^2 = 0$, due to graded-commutativity. Set

$$(3.1) \quad \mathcal{R}_d^i(A^\bullet) = \{a \in A^1 \mid \dim_{\mathbb{C}} H^i(A^\bullet, \mu_a) \geq d\}.$$

Remark 3.1. It seems worth pointing out that, given an arbitrary group extension (2.2), one has the following symmetry property, related to resonance. By standard homological algebra (see e.g. [8]), the conjugation action of Γ on T induces a (right) action of D on $A^\bullet = H^\bullet(T, \mathbb{C})$, by graded algebra automorphisms. We conclude from (3.1) that the D -representation on A^1 leaves $\mathcal{R}_d^i(A^\bullet)$ invariant, for all i and d .

In this paper, we need to consider only the resonance varieties $\mathcal{R}_d(A^\bullet) := \mathcal{R}_d^1(A^\bullet)$, which plainly depend only on the co-restriction of the multiplication map, $\cup : A^1 \wedge A^1 \rightarrow A^2$, to its image. When $\dim_{\mathbb{C}} A^1 < \infty$, it is easy to see that each $\mathcal{R}_d(A^\bullet)$ is a Zariski closed, homogeneous subvariety of the affine space A^1 . For a connected space M , we denote $\mathcal{R}_d^i(H^\bullet(M, \mathbb{C}))$ by $\mathcal{R}_d^i(M)$. For a group G , $\mathcal{R}_d^i(G)$ means $\mathcal{R}_d^i(K(G, 1))$. By considering a classifying map, it is immediate to check that $\mathcal{R}_d(M) = \mathcal{R}_d(\pi_1(M))$. We will abbreviate $\mathcal{R}_1$ by $\mathcal{R}$.

There is a well-known, useful relation between cup-product in low degrees and group commutator, see Sullivan [35], and also Lambe [24] for details. For a group G with finite first Betti number, there is a short exact sequence,

$$(3.2) \quad 0 \rightarrow (\mathrm{gr}_2 G \otimes \mathbb{C})^* \xrightarrow{d} H^1 G \wedge H^1 G \xrightarrow{\cup} H^2 G,$$

where cohomology is taken with $\mathbb{C}$ -coefficients and d is dual to the Lie bracket β from (2.3). (Note that the only finiteness assumption needed in the proof from [24] is $b_1(G) < \infty$.)

In what follows, we retain the notation from Section 2. Set $V = L(\mathbb{C})^*$ and $\mathcal{R} = \mathcal{R}(T_g)$. We infer from (2.3) and (3.2) that $\mathcal{R} \subseteq V$ is $Sp_g(\mathbb{C})$ -invariant. Here is our first step in exploiting the complex symmetry from Theorem 2.3.

Lemma 3.2. *If $\mathcal{R} \neq 0$, then $\mathcal{R}$ contains a maximal vector of the $\mathfrak{sp}_g$ -module V .*

Proof. This is an easy consequence of Borel's fixed point theorem [19, Theorem 21.2], which guarantees the existence of a B -invariant line, $\mathbb{C} \cdot v \subseteq \mathcal{R}$. Invariance under the action of the maximal torus implies that v belongs to a weight space of the $\mathfrak{h}$ -action on V , that is, $v \in V_\lambda$ for some $\lambda \in \mathfrak{h}^*$. Finally, $\mathfrak{n}^+ \cdot v = 0$ follows from U -invariance. $\square$

Since -1 belongs to the Weyl group of $\mathfrak{sp}_g$, all finite-dimensional representations of $\mathfrak{sp}_g$ are self-dual [18, Exercise 6 on p.116]. This remark leads to the following dual reformulation of Theorem 2.3, via (3.2).

Lemma 3.3. *Set $V = V(\lambda_3)$. Then $H^1 T_g = V$, as $\mathfrak{sp}_g$ -modules, and the kernel of the cup-product, $\cup : \bigwedge^2 H^1 T_g \rightarrow H^2 T_g$, is $V(2\lambda_2) \oplus V(0)$.*

Our main result in this section, which proves in particular Theorem D(1), is the following.

Theorem 3.4. *For $g = 3$, $\mathcal{R}(T_3) = H^1 T_3$, while $\mathcal{R}(T_g) = 0$, for $g \geq 4$.*

The assertion for $g = 3$ is an immediate consequence of Lemma 3.3, since in this case $\bigwedge^2 V = V(2\lambda_2) \oplus V(0)$, cf. Lemma 10.2 from [16]. So, we will assume in the sequel that $g \geq 4$ and $\mathcal{R}(T_g) \neq 0$, and use Lemma 3.2 to derive a contradiction.

Set $V(0) = \mathbb{C} \cdot z_0$. We will need explicit maximal vectors, v_0 for $V(\lambda_3)$ and u_0 for $V(2\lambda_2)$. To this end, we recall that $V(\lambda_3) = (\bigwedge^3 H/H) \otimes \mathbb{C}$, where the $\mathfrak{sp}_g$ -action on $\bigwedge^\bullet H$ is by algebra derivations; in particular, $s \cdot x \wedge y \wedge z = s \cdot x \wedge y \wedge z + x \wedge s \cdot y \wedge z + x \wedge y \wedge s \cdot z$, for $s \in \mathfrak{s}$ and $x, y, z \in H$. Set $v'_0 = a_1 \wedge a_2 \wedge a_3$, and $u'_0 = \sum_{k=3}^g (a_1 \wedge a_2 \wedge a_k) \wedge (a_1 \wedge a_2 \wedge b_k)$. Denote by v_0 the class of v'_0 in $V = V(\lambda_3)$, and let u_0 be the class of u'_0 in $\bigwedge^2 V$. Using the explicit description of $\mathfrak{h} \oplus \mathfrak{n}^+$ from [16, Section 6], it is straightforward to check that v_0 has weight λ_3 , u_0 has weight $2\lambda_2$, and $\mathfrak{n}^+ \cdot v_0 = \mathfrak{n}^+ \cdot u_0 = 0$.

To verify that both v_0 and u_0 are non-zero, we will resort to the $Sp_g(\mathbb{Z})$ -equivariant contraction constructed by Johnson in [20, p.235], $C : \bigwedge^3 H \rightarrow H$, given by

$$(3.3) \quad C(x \wedge y \wedge z) = (x \cdot y)z + (y \cdot z)x + (z \cdot x)y,$$

where the dot designates the intersection form on H . Set $L' = \ker(C)$, and $L'(\mathbb{C}) = L' \otimes \mathbb{C}$. It follows from [20, pp.238-239] that there is an induced $\mathfrak{sp}_g$ -isomorphism,

$$(3.4) \quad L'(\mathbb{C}) \xrightarrow{\sim} L(\mathbb{C}).$$

Clearly, $0 \neq v'_0 \in L'$ and $0 \neq u'_0 \in \bigwedge^2 L'$. Hence, both v_0 and u_0 are non-zero, as needed.

We infer from uniqueness of maximal vectors [18, Corollary 20.2] that necessarily $v_0 \in \mathcal{R} = \mathcal{R}(T_g)$, if $\mathcal{R} \neq 0$; see Lemma 3.2. Denote by W the $\mathfrak{s}$ -submodule $V(2\lambda_2) \oplus V(0)$ of $\bigwedge^2 V$ from Lemma 3.3. By definition (3.1) and Lemma 3.3, $v_0 \in \mathcal{R}$

if and only if $\text{im}(\mu_0) \cap W \neq 0$, where $\mu_0 : V \rightarrow \bigwedge^2 V$ denotes left-multiplication by $v_0 \in V$ in the exterior algebra.

Since $\mathfrak{n}^+ \cdot v_0 = 0$, we conclude that μ_0 is $\mathfrak{n}^+$ -equivariant. Since $v_0 \in V_{\lambda_3}$, it follows that

$$(3.5) \quad \mu_0(V_{\lambda'}) \subseteq \left(\bigwedge^2 V \right)_{\lambda_3 + \lambda'},$$

for each weight subspace $V_{\lambda'} \subseteq V$.

Lemma 3.5. *If $\mathcal{R}(T_g) \neq 0$, then $\text{im}(\mu_0) \cap W$ contains a non-zero vector killed by $\mathfrak{n}^+$.*

Proof. Taking into account the preceding discussion, the assertion will follow from Engel's theorem [18, Theorem 3.3], applied to the Lie algebra $\mathfrak{n}^+$ and the $\mathfrak{n}^+$ -module $\text{im}(\mu_0) \cap W$. It is enough to check that the action of any $n \in \mathfrak{n}^+$ on W is nilpotent. This in turn follows from the fact that $s_{\alpha_1} \cdots s_{\alpha_r} \cdot W_{\lambda'} \subseteq W_{\lambda''}$, where $\lambda'' = \lambda' + \sum_{i=1}^r \alpha_i$, for any $\alpha_1, \dots, \alpha_r \in \Phi^+$ and any non-trivial weight space $W_{\lambda'}$. Indeed, $W_{\lambda''} = 0$ for r big enough. To check the vanishing claim, one may invoke [18, Theorem 20.2(b)], and then use a height argument. $\square$

Lemma 3.6. *The following hold.*

- (1) *For $g \geq 4$, the vector $v_0 \wedge u_0 \in \bigwedge^3 V$ is non-zero.*
- (2) *The class of $(a_1 \wedge a_2 \wedge a_3) \bigwedge (b_1 \wedge b_2 \wedge b_3)$ in $\bigwedge^2 V$ does not belong to $V(0)$.*

Proof. (1) By (3.4), it is enough to show that $v'_0 \wedge u'_0 \in \bigwedge^3(\bigwedge^3 H)$ is non-zero. This is clear, since $v'_0 \wedge u'_0 = \sum_{k=4}^g (a_1 \wedge a_2 \wedge a_3) \bigwedge (a_1 \wedge a_2 \wedge a_k) \bigwedge (a_1 \wedge a_2 \wedge b_k)$.

(2) Again by (3.4), it suffices to verify that the element $e = (a_1 \wedge a_2 \wedge a_3) \bigwedge (b_1 \wedge b_2 \wedge b_3) \in \bigwedge^2(\bigwedge^3 H) \otimes \mathbb{C}$ is not killed by $\mathfrak{sp}_g$, since $b_1 \wedge b_2 \wedge b_3 \in L'$. Indeed, $T_1 \cdot e = (a_1 \wedge a_2 \wedge a_3) \bigwedge (a_1 \wedge b_2 \wedge b_3) \neq 0$, where $T_1 \in \mathfrak{sp}_g$ is described in [16, Section 6]. $\square$

We will finish the proof of Theorem 3.4, by showing that Lemmas 3.5 and 3.6 lead to a contradiction, assuming $g \geq 4$ and $\mathcal{R}(T_g) \neq 0$.

Indeed, in this case there is $v \in V(\lambda_3)$ such that

$$(3.6) \quad 0 \neq v_0 \wedge v = w_0 + w,$$

with $w_0 \in V(0)$, $w \in V(2\lambda_2)$ and $\mathfrak{n}^+ \cdot w = 0$, by Lemma 3.5.

If $w \neq 0$, then $w \in \mathbb{C}^* \cdot u_0$, by uniqueness of maximal vectors. Taking the weight decomposition of v , it follows from (3.5) that we may suppose that $w_0 = 0$ in (3.6). Therefore, $v_0 \wedge u_0 = 0 \in \bigwedge^3 V$, contradicting Lemma 3.6(1).

If $w = 0$, then $v_0 \wedge v \in \mathbb{C}^* \cdot z_0$, for some $v \in V(\lambda_3)_{-\lambda_3}$, by the same argument on weight decomposition as before. Since the weights λ_3 and $-\lambda_3$ are conjugate under the action of the Weyl group, the weight space $V(\lambda_3)_{-\lambda_3}$ is one-dimensional,

generated by the class $\bar{v}_0$ of $b_1 \wedge b_2 \wedge b_3 \in L'$ in V ; see [18, Theorems 20.2(c) and 21.2], [16, Section 6] and (3.4). Therefore, $v_0 \wedge \bar{v}_0 \in V(0)$, contradicting Lemma 3.6(2). The proof of Theorem 3.4 is thus completed.

4. RESONANCE AND FINITENESS PROPERTIES

We devote this section to a general discussion of finiteness properties related to resonance, with an application to Torelli groups. Unless otherwise mentioned, we work with $\mathbb{C}$ -coefficients. For a graded object $\mathcal{O}_\bullet$, the notation $\mathcal{O}$ means that we forget the grading.

We need to review a couple of key notions. We begin with the *holonomy Lie algebra* associated to a connected CW-complex M with finite 1-skeleton, $\mathcal{H}_\bullet(M)$. This is a quadratic graded Lie algebra, defined as the quotient of the free Lie algebra generated by H_1M , $\mathbb{L}_\bullet(H_1M)$, graded by bracket length, modulo the ideal generated by the image of the comultiplication map, $\partial_M : H_2M \rightarrow \bigwedge^2 H_1M$; here we are using the standard identification $\bigwedge^2 H_1M \equiv \mathbb{L}_2(H_1M)$, given by the Lie bracket. Note that the dual of ∂_M is the cup-product, $\cup_M : \bigwedge^2 H^1M \rightarrow H^2M$.

When G is a finitely generated group and $M = K(G, 1)$, $\mathcal{H}_\bullet(M)$ is denoted $\mathcal{H}_\bullet(G)$. By considering a classifying map, it is easy to see that $\mathcal{H}_\bullet(M) = \mathcal{H}_\bullet(\pi_1(M))$, for a connected CW-complex M with finite 1-skeleton. The associated graded Lie algebra $\text{gr}_\bullet G \otimes \mathbb{C}$, denoted $\mathcal{G}_\bullet(G)$, is also finitely generated in degree one, but not necessarily quadratic. The canonical identification, $H_1G \equiv \mathcal{G}_1(G)$, extends to a graded Lie algebra epimorphism, $\mathbb{L}_\bullet(H_1G) \twoheadrightarrow \mathcal{G}_\bullet(G)$. By (3.2), this factors to a graded Lie algebra surjection,

$$(4.1) \quad \mathcal{H}_\bullet(G) \twoheadrightarrow \mathcal{G}_\bullet(G).$$

Let $P_\bullet$ be the polynomial algebra $\text{Sym}(H_1G)$, endowed with the usual grading. For a Lie algebra $\mathcal{H}$, denote by $\mathcal{H}' := [\mathcal{H}, \mathcal{H}]$ the derived Lie algebra, and by $\mathcal{H}'' := [\mathcal{H}', \mathcal{H}']$ the second derived Lie algebra. The exact sequence of graded Lie algebras

$$(4.2) \quad 0 \rightarrow \mathcal{H}'_\bullet(G)/\mathcal{H}''_\bullet(G) \rightarrow \mathcal{H}_\bullet(G)/\mathcal{H}''_\bullet(G) \rightarrow \mathcal{H}_\bullet(G)/\mathcal{H}'_\bullet(G) \rightarrow 0$$

yields a positively graded $P_\bullet$ -module structure on the *infinitesimal Alexander invariant*, $\mathfrak{b}_\bullet(G) := \mathcal{H}'_\bullet(G)/\mathcal{H}''_\bullet(G)$, induced by the adjoint action. A finite $P_\bullet$ -presentation of $\mathfrak{b}_\bullet(G)$ is described in [30, Theorem 6.2].

Let $E_{k-1}(\mathfrak{b}(G)) \subseteq P$ be the *elementary ideal* generated by the codimension $k-1$ minors of a finite P -presentation for $\mathfrak{b}(G)$. Denote by $\mathcal{W}_k(G) \subseteq H^1G$ the zero set of $E_{k-1}(\mathfrak{b}(G))$. The next lemma explains the relationship between the infinitesimal Alexander invariant and the resonance varieties in degree one.

Lemma 4.1. *Let G be a finitely generated group. Then the equality*

$$\mathcal{W}_k(G) \setminus \{0\} = \mathcal{R}_k(G) \setminus \{0\}$$

holds for all $k \geq 1$.

Proof. We know from [30, Theorem 6.2] that $\mathfrak{b}(G) = \text{coker}(\nabla)$, as P -modules, where

$$(4.3) \quad \nabla := \delta_3 + \text{id} \otimes \partial_G: P \otimes \left(\bigwedge^3 H_1 G \oplus H_2 G \right) \rightarrow P \otimes \bigwedge^2 H_1 G,$$

and the P -linear map δ_3 is given by $\delta_3(a \wedge b \wedge c) = a \otimes b \wedge c + b \otimes c \wedge a + c \otimes a \wedge b$, for $a, b, c \in H_1 G$.

Pick any $z \in H^1 G \setminus \{0\}$. By linear algebra, we infer that $z \in \mathcal{W}_k(G)$ if and only if $\dim_{\mathbb{C}} \text{coker}(\nabla(z)) \geq k$. Consider the exact cochain complex $(\bigwedge^{\bullet} H^1 G, \lambda_z)$, where λ_z denotes left multiplication by z , and the dual exact chain complex, $(\bigwedge^{\bullet} H_1 G, \sharp \lambda_z)$. It is straightforward to check that the restriction of $\sharp \lambda_z$ to $\bigwedge^3 H_1 G$ equals $\delta_3(z)$, where δ_3 is as in (4.3).

Denoting by $\delta_2(z)$ the restriction of $\sharp \lambda_z$ to $\bigwedge^2 H_1 G$, we obtain from exactness the following isomorphism:

$$\text{coker}(\nabla(z)) \cong \text{im}(\delta_2(z)) / \text{im}(\delta_2(z) \circ \partial_G).$$

By exactness again, $\dim_{\mathbb{C}} \text{im}(\delta_2(z)) = n - 1$, where $n := b_1(G)$. Hence $z \in \mathcal{W}_k(G)$ if and only if $\text{rank}(\delta_2(z) \circ \partial_G) \leq n - 1 - k$.

Plainly, the linear map dual to $\delta_2(z) \circ \partial_G$ is μ_z . Consequently, $z \in \mathcal{W}_k(G)$ if and only if $\text{rank}(\mu_z) \leq n - 1 - k$, that is, if and only if $z \in \mathcal{R}_k(G)$; see definition (3.1). $\square$

We may spell out our first general result relating resonance and finiteness.

Theorem 4.2. *Let G be a finitely generated group. Then $\mathcal{R}_1^1(G) \subseteq \{0\}$ if and only if $\dim_{\mathbb{C}} \mathfrak{b}(G) < \infty$.*

Proof. Lemma 4.1 yields in particular the equality $\mathcal{R}_1^1(G) = \text{Zero}(\text{ann } \mathfrak{b}(G))$, away from the origin. Let $\mathfrak{m} \subseteq P$ be the maximal ideal of $0 \in H^1 G$.

If $\dim_{\mathbb{C}} \mathfrak{b}(G) < \infty$, then $\mathfrak{m}^k \subseteq \text{ann } \mathfrak{b}(G)$, for some k , by degree inspection. Taking zero sets, we obtain that $\mathcal{R}_1^1(G) \subseteq \{0\}$.

Conversely, the assumption $\mathcal{R}_1^1(G) \subseteq \{0\}$ implies $\mathfrak{m} \subseteq \sqrt{\text{ann } \mathfrak{b}(G)}$, by Hilbert's Nullstellensatz. Therefore, $\mathfrak{m}^k \subseteq \text{ann } \mathfrak{b}(G)$, for some k . Since $\mathfrak{b}(G)$ is finitely generated over P , we infer that $\dim_{\mathbb{C}} \mathfrak{b}(G) < \infty$. $\square$

Theorem 4.2 has several interesting consequences. To describe them, we first recall a couple of notions from rational homotopy theory. A *Malcev Lie algebra* (in the sense of Quillen) is a Lie algebra endowed with a decreasing vector space filtration satisfying certain axioms; see [32, Appendix A], where Quillen associates in a functorial way a Malcev Lie algebra, $\mathcal{M}(G)$, to an arbitrary group G . Following Sullivan [36], we say that a finitely generated group G is *1-formal* if $\mathcal{M}(G)$ is the completion with respect to degree of a quadratic Lie algebra, as a filtered Lie algebra. In Theorem 1.1 from [16], Hain proves that the Torelli group T_g is 1-formal, for $g \geq 6$

(but T_3 is not 1-formal). Fundamental groups of compact Kahler manifolds are 1-formal, as shown by Deligne, Griffiths, Morgan and Sullivan in [10]. Many other interesting examples of 1-formal groups are known; see e.g. [13] and the references therein.

Next, we review the *Alexander invariant*, $B(G) = G'_{\text{ab}} \otimes \mathbb{C}$, associated to a finitely generated group G . The exact sequence of groups

$$(4.4) \quad 1 \rightarrow G'/G'' \rightarrow G/G'' \rightarrow G/G' \rightarrow 1$$

may be used to put a finitely generated module structure on $B(G)$, induced by conjugation, over the Noetherian group ring $\mathbb{C}G_{\text{ab}}$. We denote by $I \subseteq \mathbb{C}G_{\text{ab}}$ the augmentation ideal, and by $\widehat{B}$ the I -adic completion of a $\mathbb{C}G_{\text{ab}}$ -module B .

Given a graded vector space $\mathcal{V}_\bullet$, $\widehat{\mathcal{V}}_\bullet$ means completion with respect to the degree filtration: $\widehat{\mathcal{V}}_\bullet = \varprojlim_q \mathcal{V}/F_q$, where $F_q = \mathcal{V}_{\geq q}$. The canonical filtration of $\widehat{\mathcal{V}}_\bullet$ is $\widehat{F}_q = \ker(\pi_q)$, where $\pi_q : \widehat{\mathcal{V}}_\bullet \rightarrow \mathcal{V}/F_q$ is the q -th projection of the inverse limit. The next corollary proves Theorem A(1).

Corollary 4.3. *Let G be a finitely generated, 1-formal group. Then $\mathcal{R}_1^1(G) \subseteq \{0\}$ if and only if $\dim_{\mathbb{C}} \widehat{B(G)} < \infty$. In particular, $\mathcal{R}_1^1(G) \subseteq \{0\}$, when $\dim_{\mathbb{C}} B(G) < \infty$.*

Proof. Let $\mathfrak{m} \subseteq P$ be the maximal ideal of the origin. The 1-formality of G provides a vector space isomorphism between $\widehat{B(G)}$ and the $\mathfrak{m}$ -adic completion of $\mathfrak{b}(G)$, cf. Theorem 5.6 from [13]. Since $\mathfrak{b}_\bullet(G)$ is generated in degree 0, its $\mathfrak{m}$ -adic completion coincides with the degree completion, $\widehat{\mathfrak{b}_\bullet(G)}$. Clearly, $\mathfrak{b}_\bullet(G)$ and $\widehat{\mathfrak{b}_\bullet(G)}$ are simultaneously finite-dimensional. Hence, our first claim follows from Theorem 4.2. For the second claim, simply note that the completion of a finite-dimensional vector space is again finite-dimensional. $\square$

Remark 4.4. The finitely generated group G discussed in Example 6.4 from [31] satisfies $\mathcal{R}_1^1(G) \subseteq \{0\}$, yet $\dim_{\mathbb{C}} B(G) = \infty$. It can be shown that G is 1-formal. Consequently, $\mathcal{R}_1^1(G) \subseteq \{0\}$ is only a necessary condition for the finite-dimensionality of $B(G)$, in general.

In the particular case when G is nilpotent, the condition $\dim_{\mathbb{C}} B(G) < \infty$ is automatically satisfied, since G' is finitely generated. We recover in this way from Corollary 4.3 the resonance obstruction to 1-formality of finitely generated nilpotent groups found by Carlson and Toledo in [9, Lemma 2.4]. We refer the reader to Macinic [27], for similar higher-degree obstructions to the formality of a finitely generated nilpotent group.

As we shall see below, the main point in Corollary 4.3 is the fact that the finite-dimensionality of $\widehat{B(G)}$ forces $\mathcal{R}_1^1(G) \subseteq \{0\}$, when G is 1-formal. We point out that 1-formality is needed for this implication. Indeed, let G be the finitely generated nilpotent Heisenberg group with $b_1(G) = 2$. As noted before, $\dim_{\mathbb{C}} \widehat{B(G)} < \infty$. Yet,

the resonance variety $\mathcal{R}_1^1(G) = \mathbb{C}^2$ is non-trivial; see for instance [27, Proposition 5.5].

Let G be an arbitrary finitely generated group, and $K \subseteq G$ a subgroup containing G' . Clearly, K is normal in G , and G -conjugation makes H_1K a finitely generated module over the Noetherian group ring $\mathbb{C}[G/K]$. By restriction via the canonical epimorphism, $G/G' \twoheadrightarrow G/K$, H_1K becomes a finitely generated $\mathbb{C}G_{\text{ab}}$ -module. We may now state our next main result from this section, which proves Theorem A(2).

Theorem 4.5. *Let G be a finitely generated group, and $K \subseteq G$ a subgroup containing G' . If $\mathcal{R}_1^1(G) \subseteq \{0\}$, the vector space $\widehat{H_1K}$ is finite-dimensional.*

Proof. We first treat the particular case $K = G'$, where we know from Theorem 4.2 that the vector space $\mathcal{H}_\bullet(G)/\mathcal{H}_\bullet''(G)$ is finite-dimensional. The canonical graded Lie algebra surjection, $\mathcal{G}_\bullet(G) \twoheadrightarrow \mathcal{G}_\bullet(G/G'')$, composed with the epimorphism (4.1), gives a graded Lie algebra surjection, $\mathcal{H}_\bullet(G) \twoheadrightarrow \mathcal{G}_\bullet(G/G'')$, that factors through an epimorphism

$$(4.5) \quad \mathcal{H}_\bullet(G)/\mathcal{H}_\bullet''(G) \twoheadrightarrow \mathcal{G}_\bullet(G/G'').$$

It follows from (4.5) that $\mathcal{G}_\bullet(G/G'') = 0$, for $\bullet \gg 0$.

On the other hand, we have an isomorphism,

$$(4.6) \quad I^q \cdot H_1G' / I^{q+1} \cdot H_1G' \simeq \mathcal{G}_{q+2}(G/G''),$$

for $q \geq 0$; see [28, pp.400–401]. We infer from (4.6) that the I -adic filtration of H_1G' stabilizes, for $q \gg 0$. Therefore, $\dim_{\mathbb{C}} \widehat{H_1G'} < \infty$, as asserted.

For the general case, consider the extension

$$(4.7) \quad 1 \rightarrow G' \rightarrow K \rightarrow A \rightarrow 1,$$

where $A = K/G'$ is a finitely generated subgroup of G_{ab} , and denote by $(H_1G')_A$ the co-invariants of the A -module H_1G' , noting that the canonical projection, $H_1G' \twoheadrightarrow (H_1G')_A$, is $\mathbb{C}G_{\text{ab}}$ -linear.

The Hochschild-Serre spectral sequence of (4.7) with trivial $\mathbb{C}$ -coefficients (see e.g. [8, p.171]) provides an exact sequence of finitely generated $\mathbb{C}G_{\text{ab}}$ -modules,

$$(4.8) \quad (H_1G')_A \rightarrow H_1K \rightarrow H_1A \rightarrow 0.$$

By standard commutative algebra (see for instance [2, Chapter 10]), the I -adic completion of (4.8) is again exact. Since $\dim_{\mathbb{C}} H_1A < \infty$, our claim about $\widehat{H_1K}$ follows from the finite-dimensionality of $\widehat{H_1G'}$. $\square$

The next corollary proves Theorem D(5), thanks to Theorem 3.4.

Corollary 4.6. *Let K be a subgroup of T_g containing T'_g . The I -adic completion of $H_1(K, \mathbb{C})$ is finite-dimensional, for $g \geq 4$, where $I \subseteq \mathbb{C}[(T_g)_{\text{ab}}]$ is the augmentation ideal of the group ring.*

Denote by $K_g \subseteq T_g$ the subgroup generated by the mapping classes of Dehn twists on simple closed curves bounding in the surface Σ_g , for $g \geq 3$. Another basic result of Johnson, proved in [22], is that $K_g = \ker(J)$, where $J : T_g \rightarrow L$ is the Johnson homomorphism. We will improve Corollary 4.6 in the next section, when K contains K_g .

5. CHARACTERISTIC VARIETIES OF TORELLI GROUPS

Guided by the interplay between arithmetic and Torelli groups, coming from (2.1), we will examine now groups whose characteristic varieties possess a natural discrete symmetry. We will compute the (restricted) characteristic varieties of T_g , in degree 1, when $g \geq 4$, and deduce that $b_1(K_g) < \infty$.

Our setup in this section is the following. Let

$$(5.1) \quad 1 \rightarrow T \rightarrow \Gamma \xrightarrow{p} D \rightarrow 1$$

be a group extension, where T is finitely generated and D is an arithmetic subgroup of a complex linear algebraic group S , defined over $\mathbb{Q}$, simple and with $\mathbb{Q}$ -rank at least 1. The motivating examples are the extensions (2.1), for $g \geq 3$. Under the above assumptions on D , we recall that any finite index subgroup $D_1 \subseteq D$ is Zariski dense in S , as follows from Borel's density theorem; see e.g. [33, Corollary 5.16].

We continue by reviewing a couple of relevant facts related to character tori and characteristic varieties. Let G be a finitely generated group. The *character torus* $\mathbb{T}(G) = \text{Hom}(G_{\text{ab}}, \mathbb{C}^*)$ is a linear algebraic group, with coordinate ring the group algebra $\mathbb{C}G_{\text{ab}}$. The connected component of $1 \in \mathbb{T}(G)$ is $\mathbb{T}^0(G) = \mathbb{T}(G_{\text{abf}})$.

For the beginning, we need to assume in (5.1) only that T is finitely generated. The natural D -representation in T_{ab} (respectively T_{abf}) induced by conjugation canonically extends to $\mathbb{Z}T_{\text{ab}}$ and $\mathbb{C}T_{\text{ab}}$ (respectively to $\mathbb{Z}T_{\text{abf}}$ and $\mathbb{C}T_{\text{abf}}$). The corresponding left D -action on $\mathbb{T}(T)$, by algebraic group automorphisms, is denoted by $d \cdot \rho$, for $d \in D$ and $\rho \in \mathbb{T}(T)$, and is defined by $d \cdot \rho(u) = \rho \circ d^{-1}(u)$, for $u \in T_{\text{ab}}$.

Let now M be a connected CW-complex, with finite 1-skeleton and fundamental group $G := \pi_1(M)$. The *characteristic varieties* $\mathcal{V}_k^i(M)$ are defined for (degree) $i \geq 0$ and (depth) $k \geq 1$ by

$$(5.2) \quad \mathcal{V}_k^i(M) = \{\rho \in \mathbb{T}(G) \mid \dim_{\mathbb{C}} H_i(M, \mathbb{C}_\rho) \geq k\}.$$

Here $\mathbb{C}_\rho$ denotes the rank one complex local system on M given by the change of rings $\mathbb{Z}G \rightarrow \mathbb{C}$, corresponding to ρ . When G is a finitely generated group and $M = K(G, 1)$, we use the notation $\mathcal{V}_k^i(G)$.

If the complex M has finite q -skeleton $M^{(q)}$, for some $q \geq 1$, it is easy to check that $\mathcal{V}_k^i(M)$ is Zariski closed in $\mathbb{T}(G)$, for all $i \leq q$ and $k \geq 1$. It is equally easy to see that $\mathcal{V}_k^i(M)$ depends only on $M^{(q+1)}$, for $i \leq q$ and $k \geq 1$, when M has finite 1-skeleton and $q \geq 1$. In particular, $\mathcal{V}_k^1(M) = \mathcal{V}_k^1(\pi_1(M))$, for all $k \geq 1$. Likewise,

$\mathcal{R}_k^i(M)$ depends only on $M^{(q+1)}$, where $q \geq 1$, in the same range as before, for any M with $M^{(1)}$ finite. We will be mainly interested in the characteristic variety $\mathcal{V}(G) := \mathcal{V}_1^1(G)$ associated to a finitely generated group G .

Our starting point in this section is the following.

Lemma 5.1. *Assume in (5.1) that the group T is finitely generated. Then the characteristic varieties $\mathcal{V}_k^i(T) \subseteq \mathbb{T}(T_{\text{ab}})$ are D -invariant subsets, for all $i \geq 0$ and $k \geq 1$.*

Proof. For $\gamma \in \Gamma$, denote γ -conjugation by $\iota_\gamma : T \xrightarrow{\sim} T$. For a ring R , with group of units $R^\times$, and a group homomorphism, $\chi : T \rightarrow R^\times$, the notation R_χ means the $\mathbb{Z}T$ -module R associated to the change of rings $\chi : \mathbb{Z}T \rightarrow R$. Note that the pair $(\iota_\gamma, \text{id}_R) : (T, R_{\chi \circ \iota_\gamma}) \xrightarrow{\sim} (T, R_\chi)$, gives an isomorphism in the category of local systems; see e.g. [8, III.8]. Hence, there is an induced isomorphism, $H_*(T, R_{\chi \circ \iota_\gamma}) \xrightarrow{\sim} H_*(T, R_\chi)$. Our claim follows by taking $R = \mathbb{C}$, and inspecting definition (5.2). $\square$

Consider $\mathcal{V}_k^1(T) \cap \mathbb{T}^0(T_{\text{ab}})$. This leads us to look at a discrete group D acting linearly on a free, finitely generated abelian group L , and examine the D -invariant, Zariski closed subsets W of $\mathbb{T}(L) = \text{Hom}(L, \mathbb{C}^*)$. Besides the trivial case $W = \mathbb{T}(L)$, we find a lot of 0-dimensional examples, by taking W to be the subgroup of m -torsion elements of $\mathbb{T}(L)$. This raises a natural question: is there anything else?

To present a first answer, we need the following notions. A *translated subgroup (torus)* is a subset of the character torus $\mathbb{T} = \mathbb{T}(G)$ of the form $t \cdot \mathbb{S}$, where $t \in \mathbb{T}$ and $\mathbb{S} \subseteq \mathbb{T}$ is a closed (connected) algebraic subgroup. Note that the *direction* of the translated subgroup, $\mathbb{S}$, is uniquely determined by $t \cdot \mathbb{S}$. A Zariski closed subset $W \subseteq \mathbb{T}$ is a *union of translated tori* if each irreducible component of W is a translated torus. A fundamental result, due to Arapura [1], is that $\mathcal{V}(G)$ is a union of translated tori in $\mathbb{T}(G)$, whenever G is a *quasi-Kähler group*, that is, the fundamental group of the complement of a divisor with normal crossings in a compact Kähler manifold.

Lemma 5.2. *Let L be a D -module which is finitely generated and free as an abelian group. Assume that D is an arithmetic subgroup of a simple $\mathbb{C}$ -linear algebraic group S defined over $\mathbb{Q}$, with $\mathbb{Q} - \text{rank}(S) \geq 1$. Suppose also that the D -action on L extends to an irreducible, rational S -representation on $L(\mathbb{C}) := L \otimes \mathbb{C}$. Let $W \subset \mathbb{T}(L)$ be a D -invariant, Zariski closed, proper subset of $\mathbb{T}(L)$. If W is a union of translated tori, then W is finite.*

Proof. We know that each irreducible component of W is of the form $t \cdot \mathbb{S}$, as above. We have to show that $\dim(\mathbb{S}) = 0$. To this end, we consider the isotropy group of $t \cdot \mathbb{S}$, denoted D_1 ; it is a finite index subgroup of D . Note that $\dim(\mathbb{S}) < \dim(\mathbb{T}(L))$, since $W \neq \mathbb{T}(L)$.

The D_1 -invariance of $t \cdot \mathbb{S}$ forces the direction $\mathbb{S}$ to be D_1 -invariant as well. Therefore, the Lie algebra $T_1\mathbb{S}$ is a D_1 -invariant linear subspace of $\text{Hom}(L, \mathbb{C}) = L(\mathbb{C})^*$. Since $D_1 \subseteq S$ is Zariski dense, it follows that the subspace $T_1\mathbb{S}$ is actually S -invariant. Hence, $T_1\mathbb{S} = 0$, due to S -irreducibility. $\square$

To improve Lemma 5.2, we need a preliminary result.

Lemma 5.3. *Let L be a D -module which is finitely generated and free as an abelian group. Then the subgroup $\mathcal{O}_t$ of $\mathbb{T}(L)$, generated by the D -orbit $D \cdot t$, is finitely generated, for any $t \in \mathbb{T}(L)$.*

Proof. Pick a $\mathbb{Z}$ -basis $\{e_i\}$ of L ; identify L with $\mathbb{Z}^n$, and $\mathbb{T}(L)$ with $(\mathbb{C}^*)^n$. For $t = (t_1, \dots, t_n) \in \mathbb{T}(L)$ and $w = \sum_i w_i e_i \in L$, set $t^w := \prod_i t_i^{w_i} \in \mathbb{C}^*$.

The elements of $D \cdot t$ are of the form $(t^{v_1}, \dots, t^{v_n})$, with $v_i = \sum_j v_{ij} e_j \in L$. Define $u_k^{ij} = \delta_{ik} e_j \in L$, for $1 \leq i, j, k \leq n$. The equality

$$(t^{v_1}, \dots, t^{v_n}) = \prod_{1 \leq i, j \leq n} (t^{u_1^{ij}}, \dots, t^{u_n^{ij}})^{v_{ij}}$$

gives the desired conclusion. $\square$

Now, we are going to show that the D -irreducibility of L is inherited by the affine torus $\mathbb{T}(L)$, in the sense explained below. This proves Theorem B.

Theorem 5.4. *Let L be a D -module which is finitely generated and free as an abelian group. Assume that D is an arithmetic subgroup of a simple $\mathbb{C}$ -linear algebraic group S defined over $\mathbb{Q}$, with $\mathbb{Q}$ -rank(S) ≥ 1 . Suppose also that the D -action on L extends to an irreducible, rational S -representation on $L(\mathbb{C}) := L \otimes \mathbb{C}$. Let $W \subset \mathbb{T}(L)$ be a D -invariant, Zariski closed, proper subset of $\mathbb{T}(L)$. Then W is finite.*

Proof. We first claim that $D \cdot t$ is finite, for every $t \in W$. This is a consequence of a conjecture of Lang in diophantine geometry, proved by Laurent in [25]. We will apply Laurent's results to the closed subvariety $W \subseteq \mathbb{T}(L)$ and the subgroup $\mathcal{O}_t \subseteq \mathbb{T}(L)$, which is finitely generated, see Lemma 5.3. By [25, Théorème 2], we have an inclusion

$$(5.3) \quad W \cap \mathcal{O}_t \subseteq \bigcup \gamma \cdot \mathbb{S},$$

where the translated subgroup $\gamma \cdot \mathbb{S}$ has origin $\gamma \in \mathcal{O}_t$, is contained in W , and $\gamma \cdot \mathbb{S}$ is maximal with these properties. Moreover, there are finitely many such maximal translated subgroups, according to [25, Lemme 4].

Denote by $V \subseteq W$ the right-hand side of (5.3). Since $\mathcal{O}_t$ is D -invariant, by construction, it follows that V is D -invariant as well. Hence, V must be finite, by Lemma 5.2. Since clearly $D \cdot t \subseteq W \cap \mathcal{O}_t$, we obtain our claim.

Supposing that W is infinite, we may find a smooth point $t \in W$ whose tangent space satisfies $0 \neq T_t W \neq T_t \mathbb{T}(L)$. Translating to the origin, we obtain a D_t -invariant, proper and non-trivial linear subspace of $L(\mathbb{C})^*$, where D_t is the isotropy group of t . By the first step of our proof, D_t has finite index in D , which contradicts the S -irreducibility of $L(\mathbb{C})$. $\square$

We go on by establishing a relation between characteristic and resonance varieties in degree 1, for finitely generated groups.

Lemma 5.5. *Let G be a finitely generated group.*

- (1) *There is a finitely presented group $\overline{G}$, together with a group epimorphism, $\phi : \overline{G} \twoheadrightarrow G$, such that $\phi_{\text{ab}} : \overline{G}_{\text{ab}} \xrightarrow{\cong} G_{\text{ab}}$ is an isomorphism, inducing identifications, $\mathcal{V}_k^1(\overline{G}) \equiv \mathcal{V}_k^1(G)$ and $\mathcal{R}_k^1(\overline{G}) \equiv \mathcal{R}_k^1(G)$, for all $k \geq 1$.*
- (2) *In particular, the tangent cone at 1 of $\mathcal{V}_k^1(G)$, $TC_1 \mathcal{V}_k^1(G)$, is contained in $\mathcal{R}_k^1(G)$, for all $k \geq 1$.*

Proof. Part (1). Let X be a classifying space for G , having 1-skeleton equal to a finite wedge of circles. Denote by $\{\mathbb{Z}G_{\text{ab}} \otimes C_{\bullet} X \xrightarrow{D_{\bullet}} \mathbb{Z}G_{\text{ab}} \otimes C_{\bullet-1} X\}$ the cellular chain complex of the universal abelian cover X^{ab} , where $\{C_{\bullet} X \xrightarrow{d_{\bullet}} C_{\bullet-1} X\}$ is the cellular chain complex of X , over $\mathbb{Z}$. Since the ring $\mathbb{Z}G_{\text{ab}}$ is noetherian, $\text{im}(D_2)$ is generated over $\mathbb{Z}G_{\text{ab}}$ by the images of finitely many 2-cells, say $\{e_1, \dots, e_r\}$.

Consider now the comultiplication map, $\partial_{X^{(2)}} : H_2(X^{(2)}, \mathbb{C}) \rightarrow \wedge^2 H_1(X^{(2)}, \mathbb{C})$. Pick finitely many 2-cells, say $\{e'_1, \dots, e'_s\}$, such that the ∂ -images of the $d_{\bullet}$ -cycles belonging to $\mathbb{Z} - \text{span}\{e'_1, \dots, e'_s\}$ generate $\text{im}(\partial_{X^{(2)}})$ over $\mathbb{C}$.

Let Y be the finite subcomplex of $X^{(2)}$ obtained from $X^{(1)}$ by attaching the cells $\{e_i\}$ and $\{e'_j\}$. Set $\overline{G} := \pi_1(Y)$. Attach cells of dimension at least 3 to Y , in order to obtain a classifying space for $\overline{G}$, denoted $\overline{X}$. Extend the inclusion $Y \hookrightarrow X^{(2)}$ to a map $\overline{X} \rightarrow X$, and consider the induced group epimorphism, $\phi : \overline{G} \twoheadrightarrow G$.

By our choice of the e -cells, we infer that ϕ induces in turn an isomorphism, $H_1(\overline{G}, R_{\chi \circ \phi}) \xrightarrow{\cong} H_1(G, R_{\chi})$, for any group homomorphism, $\chi : G \rightarrow R^{\times}$, when the ring R is commutative. In particular, our claims on ϕ_{ab} and characteristic varieties are thus verified. To check the claim on resonance varieties, we may replace $\overline{G}$ by Y and G by $X^{(2)}$. From our choice for the e' -cells, we deduce that, upon identifying $\wedge^2 H_1(Y, \mathbb{C})$ and $\wedge^2 H_1(X^{(2)}, \mathbb{C})$ via ϕ , we have the equality $\text{im}(\partial_Y) = \text{im}(\partial_{X^{(2)}})$. By duality, the cup-product maps $\cup_Y$ and $\cup_{X^{(2)}}$ have isomorphic co-restrictions to the image, which proves our last claim.

Part (2). Libgober proved in [26] that $TC_1 \mathcal{V}_k^i(M) \subseteq \mathcal{R}_k^i(M)$, for $i \geq 0$ and $k \geq 1$, when M is a connected finite CW-complex. This gives the desired inclusion, for G finitely presented. By Part (1), the inclusion still holds for finitely generated groups. $\square$

Here is our main result in this section, relating arithmetic symmetry and finiteness properties, which proves Theorem C.

Corollary 5.6. *Assume in (5.1) that T is finitely generated and $\mathcal{R}(T) \subseteq \{0\}$, and $D \subseteq S$ is arithmetic, where the $\mathbb{C}$ -linear algebraic group S is defined over $\mathbb{Q}$, simple, with $\mathbb{Q}$ -rank(S) ≥ 1 . Suppose moreover that the canonical D -representation in T_{abf} extends to an irreducible, rational S -representation in $T_{\text{abf}} \otimes \mathbb{C}$. Then the following hold.*

- (1) *The restricted characteristic varieties $\mathcal{V}_k^1(T) \cap \mathbb{T}^0(T)$ are finite, for all $k \geq 1$.*
- (2) *The first Betti number of $\pi^{-1}(A)$ is finite, for any subgroup $A \subseteq T_{\text{abf}}$, where $\pi : T \twoheadrightarrow T_{\text{abf}}$ is the canonical projection.*
- (3) *If moreover $b_1(T) > 1$, the Alexander polynomial Δ^T is a non-zero constant $c \in \mathbb{Z}$, modulo the units of $\mathbb{Z}T_{\text{abf}}$.*

Proof. We may clearly assume $b_1(T) \neq 0$. We want to use Theorem 5.4, applied to the D -module $L = T_{\text{abf}}$ and the closed subvariety $W = \mathcal{V}_k^1(T) \cap \mathbb{T}^0(T)$. The D -invariance of W follows from Lemma 5.1. The fact that $W \neq \mathbb{T}(L)$ is a consequence of our assumption on $\mathcal{R}(T)$, due to Lemma 5.5(2). By resorting to Theorem 5.4, we conclude the proof of Part (1).

Set $K := \ker(\pi)$. A basic result of Dwyer and Fried [14], as refined in [31, Corollary 6.2], says that the finiteness of $\mathcal{V}_1^1(T) \cap \mathbb{T}^0(T)$ is equivalent to $\dim_{\mathbb{C}}(K_{\text{ab}}) \otimes \mathbb{C} < \infty$, for any finitely generated group T . Consider now the extension $1 \rightarrow K \rightarrow \pi^{-1}(A) \rightarrow A \rightarrow 1$. A standard application of the Hochschild-Serre spectral sequence shows that the first Betti number of $\pi^{-1}(A)$ is finite, since both $b_1(K)$ and $b_1(A)$ are finite. This completes the proof of Part (2).

Part (3) follows from Part (1), via Corollary 3.2 from [12]. $\square$

Corollary 5.6 leads to the following consequences, for Torelli groups, which prove Theorem D(2)-(4). Let $K_g \subseteq T_g$ be the Johnson kernel from Section 4.

Corollary 5.7. *Assume $g \geq 4$.*

- (1) *The intersection $\mathcal{V}_k^1(T_g) \cap \mathbb{T}^0(T_g)$ is finite, for all $k \geq 1$.*
- (2) *The vector space $H_1(N, \mathbb{C})$ is finite-dimensional, for any subgroup N of T_g containing K_g .*
- (3) *The Alexander polynomial Δ^{T_g} is a non-zero constant $c \in \mathbb{Z}$, modulo units.*

Proof. The condition on resonance is guaranteed by Theorem 3.4. Since $b_1(T_g)$ is an increasing function of g , for $g \geq 3$, $b_1(T_g) \geq 14$, when $g \geq 3$. $\square$

6. SIGMA-INVARIANTS AND THE KÄHLER PROPERTY

We close by examining groups with natural arithmetic symmetry, at the level of Bieri-Neumann-Strebel-Renz invariants. We first review briefly the definitions and the main properties; for more details and references, see e.g. [31].

We start with the *Novikov-Sikorav completion* of a finitely generated group G , with respect to $\chi \in \text{Hom}(G, \mathbb{R})$, denoted $\widehat{\mathbb{Z}G}_{-\chi}$. For $k \in \mathbb{Z}$, let F_k be the abelian subgroup of $\mathbb{Z}G$ generated by the elements $g \in G$ with $\chi(g) \geq k$. The completion, $\widehat{\mathbb{Z}G}_{-\chi}$, of $\mathbb{Z}G$ with respect to the decreasing filtration $\{F_k\}_{k \in \mathbb{Z}}$ becomes in a natural way a ring, containing the group ring $\mathbb{Z}G$.

The *Sigma-invariants* $\Sigma^q(G, \mathbb{Z})$ are defined for $q \geq 1$ by

$$(6.1) \quad \Sigma^q(G, \mathbb{Z}) = \{0 \neq \chi \in H^1(G, \mathbb{R}) \mid H_{\leq q}(G, \widehat{\mathbb{Z}G}_{-\chi}) = 0\}.$$

When the group G is of type FP_k , the above definition coincides with the one introduced by Bieri and Renz in [5], for $q \leq k$. Note that property FP_1 simply means finite generation of G . The Sigma-invariant $\Sigma^1(G, \mathbb{Z})$ of a finitely generated group G , denoted $\Sigma(G)$, coincides with the one defined by Bieri, Neumann and Strebel in [4].

It turns out that $\Sigma^q(G, \mathbb{Z})$ is an open (possibly empty) conical subset of $H^1(G, \mathbb{R})$, for all $q \leq k$, when G is of type FP_k . Moreover, in this case we have the following fundamental property. Given a group epimorphism, $\nu : G \twoheadrightarrow L$, onto an abelian group, $\ker(\nu)$ is of type FP_q , with $q \leq k$, if and only if

$$(6.2) \quad \nu^*(H^1(L, \mathbb{R}) \setminus \{0\}) \subseteq \Sigma^q(G, \mathbb{Z}).$$

Here is the analog of Lemma 5.1 for Sigma-invariants.

Lemma 6.1. *Let $p : \Gamma \twoheadrightarrow D$ be a group epimorphism, with finitely generated kernel T . Then $\Sigma^q(T, \mathbb{Z})$ is invariant under the canonical action of D on $H^1(T, \mathbb{R})$, coming from (5.1), for all $q \geq 1$.*

Proof. Novikov-Sikorav completion is functorial, in the following sense. Let $\phi : G \rightarrow K$ be a group homomorphism and $\chi \in \text{Hom}(K, \mathbb{R})$. The induced ring homomorphism, $\phi : \mathbb{Z}G \rightarrow \mathbb{Z}K$, clearly preserves the defining filtrations of $\widehat{\mathbb{Z}G}_{-\chi \circ \phi}$ and $\widehat{\mathbb{Z}K}_{-\chi}$. Passing to completions, ϕ extends to a ring homomorphism, $\widehat{\phi} : \widehat{\mathbb{Z}G}_{-\chi \circ \phi} \rightarrow \widehat{\mathbb{Z}K}_{-\chi}$.

The above remark may be applied to γ -conjugation, $\phi = \iota_\gamma : T \xrightarrow{\sim} T$, for any $\gamma \in \Gamma$, and an arbitrary additive character $\chi \in H^1(T, \mathbb{R})$. The pair $(\phi, \widehat{\phi})$ gives then an isomorphism, $(T, \widehat{\mathbb{Z}T}_{-\chi \circ \phi}) \xrightarrow{\sim} (T, \widehat{\mathbb{Z}T}_{-\chi})$, in the category of local systems. Consequently, there is an induced isomorphism, $H_*(T, \widehat{\mathbb{Z}T}_{-\chi \circ \phi}) \xrightarrow{\sim} H_*(T, \widehat{\mathbb{Z}T}_{-\chi})$. We infer from (6.1) that $\chi \in \Sigma^q(T, \mathbb{Z})$ if and only if $\chi \cdot d \in \Sigma^q(T, \mathbb{Z})$, where $d = p(\gamma)$. $\square$

Remark 6.2. Note that $-\text{id} \in Sp_g(\mathbb{Z})$ acts by $-\text{id}$ on $\wedge^3 H/H$. Consequently, $-\Sigma(T_g) = \Sigma(T_g)$. This symmetry property of $\Sigma(G)$ about the origin does not hold in general.

Note also that, when $-\Sigma(G) = \Sigma(G)$, $\Sigma(G) \neq \emptyset$ if and only if there is a finitely generated, normal subgroup N of G , with infinite abelian quotient G/N . Moreover,

in this statement G/N may be replaced by $\mathbb{Z}$. Indeed, assuming G/N to be infinite abelian, we infer that $\Sigma(G) \neq \emptyset$, by resorting to (6.2). Conversely, we know that the image of $\Sigma(G)$ in the quotient sphere, $(H^1(G, \mathbb{R}) \setminus \{0\})/\mathbb{R}_+$, is open and nonvoid. The density of rational points on this sphere [4, p.451] implies then that $\Sigma(G)$ contains an epimorphism, $\nu : G \twoheadrightarrow \mathbb{Z}$. By antipodal symmetry of $\Sigma(G)$ and (6.2) again, $\ker(\nu)$ must be finitely generated.

Under additional hypotheses, Lemma 6.1 may be used to obtain strong information on finiteness properties.

Proposition 6.3. *Let $p : \Gamma \twoheadrightarrow D$ be a group epimorphism, with finitely generated kernel T . Assume that D is an arithmetic subgroup of a complex, simple linear algebraic group S , defined over $\mathbb{Q}$, with $\mathbb{Q} - \text{rank}(S) \geq 1$, and the canonical D -action on T_{abf} induced by conjugation extends to a non-trivial, irreducible rational representation of S in $T_{\text{abf}} \otimes \mathbb{C}$. Suppose moreover that*

- (1) *either $\Sigma(T)$ is a finite disjoint union of finite intersections of open half-spaces in $H^1(T, \mathbb{R})$,*
- (2) *or $\Sigma(T)$ is the complement of a finite union of linear subspaces in $H^1(T, \mathbb{R})$.*

Then the kernel K of the natural map, $T \twoheadrightarrow T_{\text{abf}}$, is finitely generated if and only if $\Sigma(T) \neq \emptyset$.

Proof. According to (6.2), finite generation of K is equivalent to $\Sigma(T) = H^1(T, \mathbb{R}) \setminus \{0\}$. Note that our irreducibility assumptions imply that $b_1(T) > 1$. We claim that if $\Sigma(T)$ is a proper, non-void subset of $H^1(T, \mathbb{R}) \setminus \{0\}$, then there is a proper, non-trivial linear subspace $E \subseteq H^1(T, \mathbb{R})$ invariant under the canonical action of a finite index subgroup $D_0 \subseteq D$. Granting the claim, we may use the fact that D_0 is Zariski dense in S to infer that $E \otimes \mathbb{C} \subseteq (T_{\text{abf}} \otimes \mathbb{C})^*$ is S -invariant, a contradiction. Thus, we only need to verify the above claim, in order to finish the proof.

(1) In this case, we know that $\Sigma(T) = \cup_{i=1}^r C_i$, where each C_i is a chamber of a non-void, finite hyperplane arrangement $\mathcal{A}_i$ in $H^1(T, \mathbb{R})$, and the union is disjoint. By Lemma 6.1, there is a finite index subgroup $D_1 \subseteq D$ such that $C_1 \cdot d = C_1$, for any $d \in D_1$. Consider the supporting hyperplanes of C_1 , that is, the set $\mathcal{S}_1$ consisting of those hyperplanes $E \subseteq H^1(T, \mathbb{R})$ with the property that the intersection of E with the boundary of C_1 has non-void interior in E . Standard arguments show that $\mathcal{S}_1$ is a non-void subset of $\mathcal{A}_1$; see [7, Chapter V.1]. Clearly, $\mathcal{S}_1$ is D_1 -invariant, so we may choose D_0 to be the isotropy group of an element $E \in \mathcal{S}_1$.

(2) In the second case, the complement of $\Sigma(T)$ is the union of a non-void, finite arrangement $\mathcal{A}$ of non-trivial, proper linear subspaces of $H^1(T, \mathbb{R})$. Again by Lemma 6.1, $\mathcal{A}$ is D -invariant. Clearly, the isotropy group D_0 of $E \in \mathcal{A}$ satisfies the desired conditions. $\square$

Property (1) above is verified by 3-manifold groups (that is, fundamental groups of compact, connected, differentiable 3-manifolds), according to [4]. Property (2)

holds for Kahler groups, as shown by Delzant in [11]. For Torelli groups, Proposition 6.3(2) may be improved to obtain Theorem E.

Corollary 6.4. *If the Torelli group T_g ($g \geq 4$) is a Kahler group, then the Johnson kernel K_g is finitely generated.*

Proof. We have to show that $\Sigma(T_g) \neq \emptyset$, for $g \geq 4$, assuming the Kahler property. According to [11], the complement of $\Sigma(T_g)$ is a finite union,

$$\bigcup_{\alpha} f_{\alpha}^* H^1(C_{\alpha}, \mathbb{R}),$$

coming from irrational pencils on the compact Kahler manifold M , if $T_g = \pi_1(M)$. More precisely, each $f_{\alpha} : M \rightarrow C_{\alpha}$ is a holomorphic map onto a smooth compact curve with $\chi(C_{\alpha}) \leq 0$, having connected fibers, and $\dim_{\mathbb{R}} f_{\alpha}^* H^1(C_{\alpha}, \mathbb{R}) = b_1(C_{\alpha})$. If $b_1(C_{\alpha}) < b_1(T_g)$, for every α , then we are done. Otherwise, $\chi(C_{\alpha}) < 0$, for some α , since $b_1(T_g) > 2$. We infer from definition (3.1) that $\mathcal{R}_1^1(C_{\alpha}) = H^1(C_{\alpha}, \mathbb{C})$ and $f_{\alpha}^* H^1(C_{\alpha}, \mathbb{R}) \subseteq \mathcal{R}_1^1(T_g) \cap H^1(T_g, \mathbb{R})$, which contradicts Theorem 3.4. $\square$

Example 6.5. We point out that there exist Kahler groups with $\mathcal{R}_1^1(G) = 0$ and $\Sigma(G) = \emptyset$. In particular, (6.2) implies that in this situation the kernel of the canonical projection, $G \twoheadrightarrow G_{\text{abf}}$, is not finitely generated. At the same time, the condition on $\mathcal{R}_1^1(G)$ implies that there is no group epimorphism, $G \twoheadrightarrow \pi_1(\Sigma_h)$, when the genus h is at least 2, by the argument in the proof of Corollary 6.4. We give such an example, inspired by a construction of Beauville [3, Example 1.8].

Let g be a fixed-point free involution of a 1-connected compact Kahler manifold E . The existence of such an object follows for instance from Serre's result [34], which guarantees the realizability of finite groups as fundamental groups of smooth, projective complex varieties. Let $\mathbb{Z}_2$ act on the Fermat curve $F := \{x^4 + y^4 + z^4 = 0\} \subseteq \mathbb{CP}^2$ by $g(x : y : z) = (y : x : z)$. Set $M := F \times E / \mathbb{Z}_2$, where the quotient is taken with respect to the diagonal action, and $C := F / \mathbb{Z}_2$. Note that M is a compact Kahler manifold, and C is an elliptic curve. Set $G := \pi_1(M)$.

The first projection induces a holomorphic surjection, $f : M \rightarrow C$, having connected fibers, and 4 multiple fibers (of multiplicity 2). By Delzant [11], the subspace $f^* H^1(C, \mathbb{R})$ is contained in the complement of $\Sigma(G)$. Since $f^{\bullet} : H^{\bullet} C \rightarrow H^{\bullet} M$ may be identified with the inclusion of fixed points, $(H^{\bullet} F)^{\mathbb{Z}_2} \hookrightarrow (H^{\bullet} F \otimes H^{\bullet} E)^{\mathbb{Z}_2}$, and $b_1(E) = 0$, we infer that f induces in cohomology an isomorphism in degree 1 and a monomorphism in degree 2. It follows that $\Sigma(G) = \emptyset$ and $\mathcal{R}_1^1(G) = 0$, as asserted.

By a similar construction, we may exhibit examples of Kahler groups with arbitrary (non-zero) even first Betti number, having the property that $\mathcal{R}_1^1(G) = 0$ and $\Sigma(G) \neq H^1(G, \mathbb{R}) \setminus \{0\}$. As before, these conditions imply that the kernel of the canonical projection, $G \twoheadrightarrow G_{\text{abf}}$, is not finitely generated, and there is no group epimorphism, $G \twoheadrightarrow \pi_1(\Sigma_h)$, for $h \geq 2$.

Remark 6.6. In the proof of Corollary 6.4, our strategy involves two steps. Firstly, group surjections $T_g \twoheadrightarrow \pi_1(\Sigma_h)$ with $h \geq 2$ are ruled out with the aid of our result on resonance, from Theorem 3.4. Secondly, we use the symplectic symmetry of Sigma-invariants to prove non-existence of group surjections onto orbifold fundamental groups in genus 1 (or, elliptic pencils with multiple fibers, in the geometric language from [11]), $T_g \twoheadrightarrow \pi_1^{\text{orb}}(\Sigma_1)$. Example 6.5 shows that the second step is needed for the proof of Corollary 6.4.

Acknowledgment. We are grateful to Alex Suciu, for useful discussions at an early stage of this work.

REFERENCES

1. D. Arapura, *Geometry of cohomology support loci for local systems I*, J. Algebraic Geom. **6** (1997), no. 3, 563–597. 5
2. M. F. Atiyah, I. G. MacDonald, *Introduction to commutative algebra*, Addison-Wesley, Reading, Massachusetts, 1969. 4
3. A. Beauville, *Annulation du H^1 pour les fibrés en droites plats*, in: *Complex algebraic varieties (Bayreuth, 1990)*, pp. 1–15, Lecture Notes in Math., vol. 1507, Springer, Berlin, 1992. 6.5
4. R. Bieri, W. Neumann, R. Strebel, *A geometric invariant of discrete groups*, Invent. Math. **90** (1987), no. 3, 451–477. 1, 6, 6.2, 6
5. R. Bieri, B. Renz, *Valuations on free resolutions and higher geometric invariants of groups*, Comment. Math. Helvetici **63** (1988), no. 3, 464–497. 1, 6
6. D. Biss, B. Farb, *$\mathcal{K}_g$ is not finitely generated*, Invent. Math. **163** (2006), no. 1, 213–226; erratum **178** (2009), no. 1, 229. 1
7. N. Bourbaki, *Groupes et algèbres de Lie, Chapitres 4–6*, Hermann, Paris, 1968. 6
8. K. S. Brown, *Cohomology of groups*, Grad. Texts in Math., vol. 87, Springer-Verlag, New York-Berlin, 1982. 3.1, 4, 5
9. J. A. Carlson, D. Toledo, *Quadratic presentations and nilpotent Kähler groups*, J. Geom. Anal. **5** (1995), no. 3, 359–377; erratum **7** (1997), no. 3, 511–514. 4.4
10. P. Deligne, P. Griffiths, J. Morgan, D. Sullivan, *Real homotopy theory of Kähler manifolds*, Invent. Math. **29** (1975), no. 3, 245–274. 1, 4
11. T. Delzant, *L’invariant de Bieri Neumann Strebel des groupes fondamentaux des variétés kählériennes*, Math. Annalen, posted on December 22, 2009, doi:10.1007/s00208-009-0468-8 (to appear in print). 1, 6, 6, 6.5, 6.6
12. A. Dimca, S. Papadima, A. Suciu, *Alexander polynomials: Essential variables and multiplicities*, Int. Math. Research Notices vol. **2008** (2008), no. 3, Art. ID rnm119, 36 pp. 5
13. A. Dimca, S. Papadima, A. Suciu, *Topology and geometry of cohomology jump loci*, Duke Math. Journal **148** (2009), no. 3, 405–457. 4, 4
14. W. G. Dwyer, D. Fried, *Homology of free abelian covers. I*, Bull. London Math. Soc. **19** (1987), no. 4, 350–352. 1, 5
15. B. Farb, *Some problems on mapping class groups and moduli space*, in: *Problems on mapping class groups and related topics*, pp. 11–55, Proc. Sympos. Pure Math., vol. 74, Amer. Math. Soc., Providence, RI, 2006. 1
16. R. Hain, *Infinitesimal presentations of the Torelli groups*, J. Amer. Math. Soc. **10** (1997), no. 3, 597–651. 1, 1, 2, 2, 3, 3, 3, 4

17. R. Hain, *Finiteness and Torelli spaces*, in: *Problems on mapping class groups and related topics*, pp. 57–70, Proc. Sympos. Pure Math., vol. 74, Amer. Math. Soc., Providence, RI, 2006. [1](#)
18. J. E. Humphreys, *Introduction to Lie algebras and representation theory*, Grad. Texts in Math., vol. 9, Springer-Verlag, New York, 1972. [2](#), [3](#), [3](#), [3](#), [3](#)
19. J. E. Humphreys, *Linear algebraic groups*, Grad. Texts in Math., vol. 21, Springer-Verlag, New York, 1975. [2](#), [3](#)
20. D. Johnson, *An abelian quotient of the mapping class group $\mathcal{T}_g$* , Math. Ann. **249** (1980), 225–242. [1](#), [2](#), [3](#), [3](#)
21. D. Johnson, *The structure of the Torelli group I: A finite set of generators for $\mathcal{T}$* , Ann. of Math. **118** (1983), 423–442. [1](#), [2](#)
22. D. Johnson, *The structure of the Torelli group II: A characterization of the group generated by twists on bounding curves*, Topology **24** (1985), no. 2, 113–126. [1](#), [2](#), [4](#)
23. D. Johnson, *The structure of the Torelli group III: The abelianization of $\mathcal{T}$* , Topology **24** (1985), no. 2, 127–144. [1](#), [1](#), [2](#)
24. L. A. Lambe, *Two exact sequences in rational homotopy theory relating cup products and commutators*, Proc. Amer. Math. Soc. **96** (1986), no. 2, 360–364. [3](#), [3](#)
25. M. Laurent, *Equations diophantiennes exponentielles*, Invent. Math. **78** (1984), no. 2, 299–327. [1](#), [5](#), [5](#)
26. A. Libgober, *First order deformations for rank one local systems with a non-vanishing cohomology*, Topology Appl. **118** (2002), no. 1-2, 159–168. [5](#)
27. A. Macinic, *Cohomology rings and formality properties of nilpotent groups*, J. Pure Appl. Algebra **214** (2010), 1818–1826. [4.4](#)
28. W. S. Massey, *Completion of link modules*, Duke Math. J. **47** (1980), no. 2, 399–420. [1](#), [4](#)
29. D. McCullough, A. Miller, *The genus 2 Torelli group is not finitely generated*, Topology Appl. **22** (1986), no. 1, 43–49. [1](#)
30. S. Papadima, A. Suciu, *Chen Lie algebras*, Int. Math. Res. Notices **2004**, no. 21, 1057–1086. [4](#), [4](#)
31. S. Papadima, A. Suciu, *Bieri–Neumann–Strebel–Renz invariants and homology jumping loci*, Proc. London Math. Soc., posted on December 22, 2009, doi:10.1112/plms/pdp045 (to appear in print). [1](#), [4.4](#), [5](#), [6](#)
32. D. Quillen, *Rational homotopy theory*, Ann. of Math. **90** (1969), 205–295. [4](#)
33. M. S. Raghunathan, *Discrete subgroups of Lie groups*, Ergebnisse der Math., vol. 68, Springer-Verlag, Berlin, 1972. [2](#), [5](#)
34. J.-P. Serre, *Sur la topologie des variétés algébriques en caractéristique p* , in: *Symposium internacional de topología algebraica* (Mexico City, 1958), pp. 24–53. [6.5](#)
35. D. Sullivan, *On the intersection ring of compact three manifolds*, Topology **14** (1975), 275–277. [3](#)
36. D. Sullivan, *Infinitesimal computations in topology*, Inst. Hautes Études Sci. Publ. Math. **47** (1977), 269–331. [1](#), [4](#)

LABORATOIRE J.A. DIEUDONNÉ, UMR DU CNRS 6621, UNIVERSITÉ DE NICE SOPHIA-ANTIPOLIS, PARC VALROSE, 06108 NICE CEDEX 02, FRANCE

E-mail address: `dimca@unice.fr`

INSTITUTE OF MATHEMATICS SIMION STOILOW, P.O. Box 1-764, RO-014700 BUCHAREST, ROMANIA

E-mail address: `Stefan.Papadima@imar.ro`